

UCLA STAT 13
**Introduction to Statistical Methods for
the Life and Health Sciences**

● **Instructor:** Ivo Dinov,
Asst. Prof. In Statistics and Neurology

● **Teaching Assistants:**
Ming Zheng, Sovia Lau, Tianwei Yu
UCLA Statistics

University of California, Los Angeles, Fall 2003
http://www.stat.ucla.edu/~dinov/courses_students.html

Stat 13, UCLA, Ivo Dinov Slide 1

UCLA STAT 13

**to just hear is to forget
to see is to remember
to do it yourself is to understand ...
(... to NOT go to class is to ... not pass ...)**

Stat 13, UCLA, Ivo Dinov Slide 2

What is Statistics? A practical example

- Michael Benton & Francisco Ayala, *Dating the Tree of Life*, Science 2003 300: 1698-1700
- **Molecular** vs. **Paleontological** dating of major branching points in the tree of life are debated
- **Molecular** date estimates are up to twice as old (due to statistical bias) as **Paleontological** dates (missing fossils).
- **Goals:** Same as that set out by Darwin: to understand *where life came from*, the *shape of evolution*, the *place of humans in nature* and to determine the *extent of modern biodiversity* and where *it is threatened*.

Stat 13, UCLA, Ivo Dinov Slide 3

What is Statistics? A practical example

- **Plants:** The first vascular land plants are found as fossils in the Silurian, and earlier evidence from possible vascular plant spores may extend the range back to the Ordovician, **475 Ma** considerably < a molecular estimate of **700 Ma**.
- **Birds:** Molecular estimates place the split of basal clades and modern orders at **70 to 120 Ma**. The oldest uncontroversial fossils of modern bird orders date from the Paleocene (**60 Ma**), much younger.
- **Mammals:** Molecular dates split of modern placentals in the mid- to Late Cretaceous (**80 to 100 Ma**). The oldest fossil representatives of modern mammals dated from the Paleocene and Eocene (**50 to 65 Ma**).

Stat 13, UCLA, Ivo Dinov Slide 4

What is Statistics? Topics!

- It is proposed that molecular dates are correct (with **confidence intervals**) and that methods exist to correct for that **error**. However, critics have pointed out several pervasive biases that make molecular dates too old.
- First, if calibration dates are too old, then all other dates **estimated** from them will also be too old.
- A second biasing factor is that undetected fast-evolving genes could **bias** estimates of timing. **Empirical and statistical studies** of vertebrate sequences suggest that such non-clock-like genes may be detected and that they do not affect **estimates** of dating. However, **statistical tests** may have low **power** and could produce consistently > dates.

Stat 13, UCLA, Ivo Dinov Slide 5

What is Statistics? A practical example

- A 3rd source of **bias** relates to polymorphism. Two species often become fixed for alternative alleles that existed as a polymorphism in their ancestral species.
- A 4th biasing **factor** is that molecular time estimates show (**skewed asymmetric distributions**, with a constrained (large numbers) younger left-end but an unconstrained (smaller numbers) older right-end.

Stat 13, UCLA, Ivo Dinov Slide 6

What is Statistics? Estimate Variation!

Data Source	Metazoa (Animals)	Bilateria (metazoans except sponges, e.g., anemones)	Deuterostomia (backboned animals)
Gene (8 G)		1200 ± 100	1001 ± 100
Protein (64 E)	930 ± 115	790 ± 60	590
Gene (4 G)	940 ± 80	700 ± 80	
Gene (18 G)		670 ± 60	600 ± 60
Gene (22 G)		830 ± 55	
Gene (50 G)	1350 ± 150 (est.)	993 ± 46	
Gene (22 G)		659 ± 131	
Protein (10 E)		627 ± 51	
Gene (MtDNA; 18S rRNA)		588 min.	586/589 min.

Stat 13, UCLA, Ivo Dinov Slide 7

What is Statistics?


- There are three main methods of probabilistic forecasting: **time-series extrapolation**; **expert judgement**; and **extrapolation of historical forecast errors**.
- Time-series** methods rely on statistical models that are fitted to historical data. These methods, however, seldom give an accurate description of the past. If many of the historical facts remain unexplained, time-series methods result in **excessively wide prediction intervals** when used for long-term forecasting.
- Expert judgement** is subjective, and **historic-extrapolation** alone may be near-sighted.

Stat 13, UCLA, Ivo Dinov Slide 16

Chapter 1: What is Statistics?

Chris Wild & George Seber
Textbook

- Polls and surveys** – we're all different; It's impossible or expensive to investigate every single person.
- Experimentation** – sample vs. population
- Observational Studies** – selection and non-response bias
- Statistics -- What is it and who uses it?
- Summary




Stat 13, UCLA, Ivo Dinov Slide 17

Newtonian science vs. chaotic science

- Article by Robert May, Nature, vol. 411, June 21, 2001
- Science we encounter at schools deals with **crisp certainties** (e.g., prediction of planetary orbits, the periodic table as a descriptor of all elements, equations describing area, volume, velocity, position, etc.)
- As soon as **uncertainty** comes in the picture it **shakes the foundation of the deterministic science**, because only **probabilistic statements** can be made in describing a phenomenon (e.g., roulette wheels, chaotic dynamic weather predictions, Geiger counter, earthquakes, etc.)
- What is then science all about** – describing absolutely certain events and laws alone, or describing more general phenomena in terms of their behavior and chance of occurring? Or may be both!

Stat 13, UCLA, Ivo Dinov Slide 18



Errors in Samples ...

- Selection bias**: Sampled population is not a representative subgroup of the population really investigated.
- Non-response bias**: If a particular subgroup of the population studied does not respond, the resulting responses may be skewed.
- Question effects**: Survey questions may be slanted or loaded to influence the result of the sampling.
- Is **quota sampling** reliable? Each interviewer is assigned a **fixed quota** of subjects (subjects district, sex, age, income exactly specified, so investigator can select those people as they liked).
- Target population** – entire group of individuals, objects, units we study.
- Study population** – a subset of the target population containing all “units” which could possibly be used in the study.
- Sampling protocol** – procedure used to select **the sample**
- Sample** – the subset of “units” about which we actually collect info.

Stat 13, UCLA, Ivo Dinov Slide 20

More terminology ...

- **Census** – attempt to sample the entire population
- **Parameter** – numerical characteristic of the population, e.g., income, age, etc. Often we want to estimate population parameters.
- **Statistic** – a numerical characteristic of the sample. (Sample) statistic is used to estimate a corresponding population parameter.
- Why do we sample at random? We draw “units” from the study population at random to avoid bias. Every subject in the study sample is equally likely to be selected. Also random-sampling allows us to calculate the likely size of the error in our sample estimates.

Slide 21 Stat 13, UCLA, Jon Dinger

More definitions ...

- How could you implement the lottery method to randomly sample 10 students from a class of 250? – list all names; assign numbers 1,2,3,...,250 to all students; Use a random-number generator to choose (10-times) a number in range [0;250]; Process students drawn.
- Random or chance error is the difference between the sample-value and the true population-value (e.g., 49% vs. 69%, in the above body-overweight example).
- Non-sampling errors (e.g., non-response bias) in the census may be considerably larger than in a comparable survey, since surveys are much smaller operations and easier to control.
- Sampling errors—arising from a decision to use a sample rather than entire population
- Unbiased procedure/protocol: (e.g., using the proportion of left-handers from a random sample to estimate the corresponding proportion in the population).
- Cluster sampling- a cluster of individuals/units are used as a sampling unit, rather than individuals.

Slide 22 Stat 13, UCLA, Jon Dinger

More terminology ...

- What are some of the non-sampling errors that plague surveys? (non-response bias, question effects, survey format effects, interviewer effects)
- If we take a random sample from one population, can we apply the results of our survey to other populations? (It depends on how similar, in the respect studied, the two populations are. In general- No! This can be a dangerous trend.)
- Are sampling households at random and interviewing people at random on the street valid ways of sampling people from an urban population? (No, since clusters (households) may not be urban in their majority.)
- Pilot surveys – after prelim investigations and designing the trial survey Q's, we need to get a “small sample” checking clearness and ambiguity of the questions, and avoid possible sampling errors (e.g., bias).

Slide 23 Stat 13, UCLA, Jon Dinger

Questions ...

- How do the following lead to biases or cause differences in response:
 - non-response
 - self-selection
 - question effects
 - survey-format effects
 - interviewer effects
 - transferring findings?

Slide 24 Stat 13, UCLA, Jon Dinger

Questions ...

- Give an example where non-representative information from a survey may be useful. Non-representative info from surveys may be used to estimate parameters of the actual sub-population which is represented by the sample. E.g., Only about 2% of dissatisfied customers complain (most just avoid using the services), these are the most-vocal reps. So, we can not make valid conclusions about the stereotype of the dissatisfied customer, but we can use this info to tract down changes in levels of complains over years.
- Why is it important to take a pilot survey?
- Give an example of an unsatisfactory question in a questionnaire. (In a telephone study: What time is it?
Do we mean Eastern/Central/Mountain/Pacific?)

Slide 25 Stat 13, UCLA, Jon Dinger

Questions ...

- Random allocation – randomly assigning treatments to units, leads to representative sample only if we have large # experimental units.
- Completely randomized design- the simplest experimental design, allows comparisons that are unbiased (not necessarily fair). Randomly allocate treatments to all experimental units, so that every treatment is applied to the same number of units. E.g., If we have 12 units and 3 treatments, and we study treatment efficacy, we randomly assign each of the 3 treatments to 4 units exactly.
- Blocking- grouping units into blocks of similar units for making treatment-effect comparisons only within individual groups. E.g., Study of human life expectancy perhaps income is clearly a factor, we can have high- and low-income blocks and compare, say, gender differences within these blocks separately.

Slide 26 Stat 13, UCLA, Jon Dinger

Questions ...

- Why should we try to “blind” the investigator in an experiment?
 - Why should we try to “blind” human experimental subjects?
 - The basic rule of experimenter :
- “Block what you can and randomize what you cannot.”

Slide 27 Stat 13, UCLA, Jon Dinger

Experiments vs. observational studies for comparing the effects of treatments

- In an Experiment
 - experimenter determines which units receive which treatments. (ideally using some form of random allocation)
- Observational study – useful when can’t design a controlled randomized study
 - compare units that happen to have received each of the treatments
 - Ideal for describing relationships between different characteristics in a population.
 - often useful for identifying possible causes of effects, but cannot reliably establish causation.
- Only properly designed and executed experiments can reliably demonstrate causation.

Slide 28 Stat 13, UCLA, Jon Dinger

The Subject of Statistics

Statistics is concerned with the process of finding out about the world and how it operates -

- in the face of variation and uncertainty
- by collecting and then making sense (interpreting) of data.

Slide 30 Stat 13, UCLA, Jon Dinger

The Role of Randomization

Well designed statistical studies employ randomization to avoid subjective and other biases.

- Surveys and observational studies should use random sampling to obtain representative samples.
- Experiments should use random assignment of experimental subjects to treatment groups
 - to ensure comparisons are fair i.e., treatment groups are as similar as possible in every way except for the treatment being used.

Slide 34 Stat 13, UCLA, Jon Dinger

“Blocking” vs. “stratification”

“Blocking”

- word used in describing an experimental design

“Stratification”

- used in describing a survey or observational study
- Both refer to idea of only making comparisons within relatively similar groups of subjects

Slide 35 Stat 13, UCLA, Jon Dinger

Blocking and randomization

“Block what you can and randomize what you cannot.”

- Block to ensure fair comparisons with respect to factors known to be important
- Randomize to try to obtain comparability with respect to unknown factors
- Randomization also allows the calculation of how much the estimates made from the study data are likely to be in error

Slide 36 Stat 13, UCLA, Jon Dinger

Sources of error in surveys

- Random sampling leads to **sampling errors**, sampling-size (as we saw for the overweight survey), arising for the choice to use a sample, as opposed to census.
- **Non-sampling errors** can be much larger than the sampling errors. Selection bias, non-response bias, survey/question/interview format are all non-sampling errors.

Slide 37 Stat 13, UCLA, Jon Dinger

Sources of non-sampling errors

- **Selection bias:**
Arises when the population sampled is not exactly the population of interest.
- **Self-selection:**
People themselves decide whether or not to be surveyed. Results akin to severe non-response.
- **Non-response bias:**
Non-respondents often behave or think differently from respondents
 - low response rates can lead to huge biases.

Slide 38 Stat 13, UCLA, Jon Dinger

Non-sampling errors cont.

- **Question-wording effects:**
Even slight differences in question wording can produce measurable differences in how people respond.
- **Interviewer effects:**
Different interviewers asking the same questions can tend to obtain different answers.
- **Survey format effects:**
Factors such as question order, questionnaire layout, self-administered questionnaire or interviewer, can effect the results.

Slide 39 Stat 13, UCLA, Jon Dinger

Dealing with errors

- **Statistical methods** are available for estimating the likely size of **sampling errors**.
- All we can do with **non-sampling errors** is to try to minimize them at the study-design stage.
- **Pilot survey:**
One tests a survey on a relatively small group of people to try to identify any problems with the survey design before conducting the survey proper.

Slide 40 Stat 13, UCLA, Jon Dinger

Jargon describing experiments

- **Control group:**
 - group of experimental units is given no treatment.
 - treatment effect estimated by comparing each treatment group with control group
- **Blinding:**
 - Preventing people involved in experiment from knowing which experimental subjects have received which treatment
 - One may be able to blind
 - subjects themselves
 - people administering the treatments
 - people measuring the results.

Slide 41 Stat 13, UCLA, Jon Dinger

Jargon describing experiments

- **Double blind:**
Both the subjects and those administering the treatments have been blinded.
- **Placebo:**
An inert/dummy/fake treatment.
- **Placebo effect:**
Response caused in human subjects by the idea that they are being treated.

Slide 42 Stat 13, UCLA, Jon Dinger

Poll Example

- A survey of High School principals taken after a widespread change in the public school system revealed that 20% of them were under stress-relief medication, and almost 50% had seen a doctor in the past 6 mo.s with stress complains. The survey was compiled from 250 questionnaires returned out of 2500 sent out. How reliable the results of this experiment are and why?

Slide 45 Stat 13, UCLA, Jon Dinger

Poll Example

- This is only a 10% response rate - the people who responded could be very unrepresentative. It could well be that the survey struck a responsive chord with stressed-out principals.

Slide 46 Stat 13, UCLA, Jon Dinger

Experimental vs. Observation study

- A researcher wants to evaluate IQ levels are related to person's height. 100 people are randomly selected and grouped into 5 bins: [0:50), [50:100), [100:150), [150:200), [200:250] cm in height. The subjects undertook a IQ exam and the results are analyzed.
- Another researcher wants to assess the bleaching effects of 10 laundry detergents on 3 different colors (R,G,B). The laundry detergents are randomly selected and applied to 10 pieces of cloth. The discoloration is finally evaluated.

Slide 47 Stat 13, UCLA, Jon Dinger

Experimental vs. Observation study

- For each study, describe what *treatment* is being compared and what *response* is being measured to compare the treatments.
- Which of the studies would be described as *experiments* and which would be described as *observational* studies?
- For the studies that are *observational*, could an experiment have been carried out instead? If not, briefly explain why not.
- For the studies that are *experiments*, briefly discuss what *forms of blinding* would be possible to be used.
- In which of the studies has *blocking* been used? Briefly describe *what* was blocked and why it was blocked.


Slide 48 Stat 13, UCLA, Jon Dinger

Experimental vs. Observation study

- What is the *treatment* and what is the *response*?
 1. *Treatment* is height (as a bin). *Response* is IQ score.
 2. *Treatment* is laundry detergent. *Response* is discoloration.
- *Experiment* or *observational* study?
 1. *Observational* – compare obs's (IQ) which happen to have the treatment (height).
 2. *Experimental* – experimenter controls which treatment is applied to which unit.
- For the *observational* studies, can we conduct an experiment?
 1. This *could not* be done as an experiment - it would require the experimenter to decide the (natural) height (treatment) of the subjects (units).
- For the *experiments*, is there *blinding*?
 2. The only form of blinding possible would be for the technicians measuring the cloth discoloration not to know which detergent was applied.
- Is there *blocking*?
 1. & 2. *No blocking*. Say, if there are two laundry machines with different cycles of operation and if we want to block we'll need to randomize which laundry does which cloth/detergent combinations, because differences in laundry cycles are a known source of variation.

Slide 49 Stat 13, UCLA, Jon Dinger


Mean, Median, Mode, Quartiles, 5# summary

- The *sample mean* is the average of all numeric obs's.
- The *sample median* is the obs. at the index $(n+1)/2$ (note take avg of the 2 obs's in the middle for fractions like 23.5), of the observations ordered by size (small-to-large)?
- The *sample median* usually preferred to the *sample mean* for *skewed data*?
 
- Under what circumstances may quoting a *single center* (be it mean or median) not make sense? (multi-modal)
- What can we say about the sample mean of a *qualitative variable*? (meaningless)

Slide 50 Stat 13, UCLA, Jon Dinger

Quartiles

The first quartile (Q_1) is the median of all the observations whose *position* is strictly below the *position* of the median, and the third quartile (Q_3) is the median of those above.



Slide 51

Stat 13, UCLA, Ivo Dinov

Five number summary

The five-number summary = (Min, Q_1 , Med, Q_3 , Max)

Slide 52

Stat 13, UCLA, Ivo Dinov