

UCLA STAT 13
**Introduction to Statistical Methods for the
 Life and Health Sciences**

Instructor: Ivo Dinov,
 Asst. Prof. of Statistics and Neurology

Teaching Assistants:
Jacqueline Dacosta & Chris Barr

University of California, Los Angeles, Fall 2006
http://www.stat.ucla.edu/~dinov/courses_students.html

Slide 1 Stat 13, UCLA, Ivo Dinov

Lecture Set 8
The T Test
Wilcoxon-Mann-Whitney Test

Slide 2 Stat 13, UCLA, Ivo Dinov

Application

Example: Nine observations of surface soil pH were made two different locations. Does the data suggest that the true mean soil pH values differ for the two locations?

Test using $\alpha = 0.05$, and be sure to check any necessary assumptions for the validity of your test.

Location 1	Location 2
8.10	7.85
7.89	7.30
8.00	7.73
7.85	7.27
8.01	7.58
7.82	7.27
7.99	7.50
7.80	7.23
7.93	7.41

Slide 3 Stat 13, UCLA, Ivo Dinov

Application

To meet the assumption of normality (necessary for the t-test with such a small sample size in each group), we will calculate a normal probability plot for each group.

Slide 4 Stat 13, UCLA, Ivo Dinov

Application

● #1 Formulate hypotheses

$H_0: \mu_1 - \mu_2 = 0$ (there is no difference between the true mean soil pH of location1 and location2)

$H_a: \mu_1 - \mu_2 \neq 0$ (there is a difference between the true mean soil pH of location1 and location2)

Slide 5 Stat 13, UCLA, Ivo Dinov

Application

● #2 Calculate the test statistic

Descriptive Statistics: Location 1, Location 2									
Variable	N	N*	Mean	SE Mean	StDev	Minimum	Q1	Median	Q3
Location 1	9	0	7.9322	0.0335	0.1005	7.8000	7.8350	7.9300	8.0050
Location 2	9	0	7.4600	0.0740	0.2220	7.2300	7.2700	7.4100	7.6550

$$SE_{\bar{y}_1 - \bar{y}_2} = \sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}} = \sqrt{\frac{0.1005^2}{9} + \frac{0.222^2}{9}} = 0.081$$


$$t_s = \frac{\bar{y}_1 - \bar{y}_2 - 0}{SE_{\bar{y}_1 - \bar{y}_2}} = \frac{7.9322 - 7.460 - 0}{0.081} = 5.827$$

Slide 6 Stat 13, UCLA, Ivo Dinov

Application

- #3 Calculate the p-value

$$df = \frac{(SE_1^2 + SE_2^2)^2}{\frac{SE_1^2}{n_1 - 1} + \frac{SE_2^2}{n_2 - 1}} = \frac{(0.0335^2 + 0.074^2)^2}{\frac{0.0335^2}{9-1} + \frac{0.074^2}{9-1}} = 11.03 \approx 11df$$



$$p < 2(0.0005) = 0.001 \text{ (SOCR)}$$

Slide 7

Stat 13, UCLA, Ivo Dinno

Application

- #4 Conclusion

Because $p < 0.001 < 0.05$, we will reject H_0 .

CONCLUSION: These data show that there [is a statistically significant true mean difference in the pH of Location 1 and Location 2](#) ($P < 0.001$).

Slide 8

Stat 13, UCLA, Ivo Dinno

Application

- Confidence interval for $\mu_1 - \mu_2$

- Suppose we calculated a 95% confidence interval to be:

$$\begin{aligned} (\bar{y}_1 - \bar{y}_2) \pm t(df)_{0.025} (SE_{\bar{y}_1 - \bar{y}_2}) &= (7.932 - 7.460) \pm t(11)_{0.025} (0.081) \\ &= (0.472) \pm 2.201(0.081) \\ &= (0.294, 0.650) \end{aligned}$$

- Does this interval surprise you?

Slide 9

Stat 13, UCLA, Ivo Dinno

Application

- Corresponding computer output:

Two-Sample T-Test and CI: Location 1, Location 2

Two-sample T for Location 1 vs Location 2

	N	Mean	StDev	SE Mean
Location 1	9	7.932	0.100	0.033
Location 2	9	7.460	0.222	0.074

Difference = mu (Location 1) - mu (Location 2)
 Estimate for difference: 0.472222
 95% CI for difference: (0.293459, 0.650985)
 T-Test of difference = 0 (vs not =): T-Value = 5.81
 P-Value = 0.000 DF = 11

Slide 10

Stat 13, UCLA, Ivo Dinno

CI and Hypothesis-Testing relationship

- Consider a 95% confidence interval for $\mu_1 - \mu_2$ and it's relationship to the t test at $\alpha = 0.05$

- Both use $\bar{y}_1 - \bar{y}_2$ and $SE_{\bar{y}_1 - \bar{y}_2}$ in their calculations

$$CI: (\bar{y}_1 - \bar{y}_2) \pm t(df)_{\alpha/2} (SE_{\bar{y}_1 - \bar{y}_2})$$

$$T_s: t_s = \frac{(\bar{y}_1 - \bar{y}_2) - 0}{SE_{\bar{y}_1 - \bar{y}_2}}$$


Slide 11

Stat 13, UCLA, Ivo Dinno

CI and Hypothesis-Testing relationship

- With a t test we reject H_0 if the p-value is less than α and fail to reject otherwise

- this is the same thing as saying we reject if t_s is beyond $\pm t_{0.025}$, and fail to reject otherwise



Slide 12

Stat 13, UCLA, Ivo Dinno

CI and Hypothesis-Testing relationship

- Focusing on the upper half of the distribution and remembering the symmetry: we fail to reject when

$$|t_s| = \frac{|\bar{y}_1 - \bar{y}_2|}{SE_{\bar{y}_1 - \bar{y}_2}} < t_{0.025}$$

- Further manipulation gives us:

$$\begin{aligned} |\bar{y}_1 - \bar{y}_2| &< t_{0.025}(SE_{\bar{y}_1 - \bar{y}_2}) \\ &= -t_{0.025}(SE_{\bar{y}_1 - \bar{y}_2}) < \bar{y}_1 - \bar{y}_2 < t_{0.025}(SE_{\bar{y}_1 - \bar{y}_2}) \\ &= -(\bar{y}_1 - \bar{y}_2) - t_{0.025}(SE_{\bar{y}_1 - \bar{y}_2}) < 0 < -(\bar{y}_1 - \bar{y}_2) + t_{0.025}(SE_{\bar{y}_1 - \bar{y}_2}) \\ &= (\bar{y}_1 - \bar{y}_2) + t_{0.025}(SE_{\bar{y}_1 - \bar{y}_2}) > 0 > (\bar{y}_1 - \bar{y}_2) - t_{0.025}(SE_{\bar{y}_1 - \bar{y}_2}) \end{aligned}$$

- Therefore, we **fail to reject H_0** : $\mu_1 - \mu_2 = 0$ (for the not equal to alternative), if the **confidence interval contains 0**.

Slide 13 Stat 13, UCLA, Ivo Dinov

CI and Hypothesis-Testing relationship

- If a two-tailed t test and a confidence interval give us the same result, why learn both?

- There are advantages to each one

- **Confidence interval**: shows magnitude of difference between μ_1 and μ_2
- **T test**: has p-value which describes the strength of evidence that μ_1 and μ_2 are really different.

Slide 14 Stat 13, UCLA, Ivo Dinov

More on the significance level α

- Choose a significance level BEFORE analyzing the data

Example: Say $df = 15$ and $\alpha = 0.05$

- If t_s is in either tail we will reject H_0 . The chance of this happening due to **random variation** is 0.05. I.e., $P(\text{reject } H_0) = 0.05$, if H_0 is true.
- Because we are assuming that H_0 is true, all t_s values on the t curve would only deviate from 0 because of **sampling error**.
- This means:
 - 95% would fail to reject H_0
 - 2.5% would reject H_0 ($-t_s$)
 - 2.5% would reject H_0 (t_s)

In other words, a total of 5% would reject H_0 when H_0 is actually true. This is an incorrect conclusion just because of sampling error!

Slide 15 Stat 13, UCLA, Ivo Dinov

More on the significance level α

- When we are analyzing one data set in real life at the 0.05 level and our conclusion is to reject H_0 there are two possible scenarios:

1. H_0 is in fact false
2. H_0 is true, but we were unlucky (5%)

Slide 16 Stat 13, UCLA, Ivo Dinov

Type I and Type II Errors

- There are two possible mistakes that can be made when conducting a hypothesis test:

- A type I error is when we reject H_0 and H_0 was true
 - **P(type I error) = α**
 - When we choose α before we conduct our test, we are actually protecting ourselves against a type I error
 - This choice will depend on your experiment
- A type II error is when we fail to reject H_0 and H_0 is false
 - **P(type II error) = β**
 - β can also be specified before we collect our data
 - will have more to do with the number of observations in our sample

Slide 17 Stat 13, UCLA, Ivo Dinov

Type I and Type II Errors

- Table (below) is the best way to summarize

		Reality	
		H_0 True	H_0 False
Decision	Fail To Reject H_0	Correct TN	Type II FN
	Reject H_0	Type I FP	Correct TP

- You cannot make both errors at the same time
 - Once you have reached a conclusion (reject or fail to reject) based on the data from your experiment you've either made a correct decision or you've made an error (type I for a reject conclusion and type II for a failing-to-reject conclusion)

Slide 18 Stat 13, UCLA, Ivo Dinov

Type I and Type II Errors

- Analogy: Think of a car with a car alarm being broken into
 - If the alarm goes off for no reason (reject H_0 when H_0 is true) - type I error
 - If the car gets broken into and the alarm doesn't go off (fail to reject H_0 when H_0 is false) - type II error
 - Also consider the sensitivity of the alarm
 - REMEMBER: Fail To Reject H_0 means "nothing is going on" or the data do not show otherwise
- Consequences of Type I / II errors are quite different

Slide 19 Stat 13, UCLA, Jon Dineen

Type I and Type II Errors

Example: Measuring pollution in a lake. Say the EPA institutes a rule that companies near bodies of water must test their pollution output. If the company doesn't find any statistical significance in their results, they may continue their current practices.

H_0 : No significant pollution

H_a : Significant pollution

In this case a type II error would be much worse (probability of failing to reject H_0 when H_0 is false – saying no significant pollution when there really is)

An "ethical" company would want to make sure they tested enough samples to guarantee that β is small

Slide 20 Stat 13, UCLA, Jon Dineen

Type I and Type II Errors

Example: Drug Treatments. Say a doctor would like to study a new highly toxic drug treatment for cancer patients. There are many risks and side effects of the new drug, but would be of benefit if the proportion of patients responding is greater than 50%.

H_0 : No significant response (Proportion responding to TX is ≤ 0.5)

H_a : Significant response (Proportion responding to TX is > 0.5)

In this case a type I error is much worse (probability of rejecting H_0 when H_0 is true – like saying that the TX does something when it really doesn't)

An ethical researcher would want to make sure they keep α small before collecting and analyzing the data

Slide 21 Stat 13, UCLA, Jon Dineen

Type I and Type II Errors

● Because α is chosen beforehand, we are protected against type I errors. However, type II errors depend on many things, such as sample size (section 7.8)

● β = P(fail to reject H_0) when H_0 is false.

■ The chance of rejecting H_0 when it is actually false is called the power of our test

Power = $1 - \beta$ = P(reject H_0) when H_0 is false

measures the ability of the test to detect a difference when a difference really does exist

Power depends on sample size. A larger sample gives more information and also increases power.

When you plan an analysis you always need to take power into account (ie plan for n):

decide desired SE and calculate n

analysis of power (7.8)

Slide 22 Stat 13, UCLA, Jon Dineen

One Tailed t Tests

● The previous hypothesis test was called a two-tailed (or non-directional) test because H_0 was rejected if t_s fell in either tail

● In some analyses it is reasonable that there will be a certain direction of a deviation from H_0

■ This means that we are looking to see if one group mean is smaller/larger than the other

● The hypotheses for a one-tailed (or directional) test are:

$H_0: \mu_1 - \mu_2 = 0$

$H_a: \mu_1 - \mu_2 > 0$

OR

$H_a: \mu_1 - \mu_2 < 0$

■ Note: the null hypothesis doesn't change

Slide 23 Stat 13, UCLA, Jon Dineen

One Tailed t Tests

● One-Tailed Test Procedure:

■ Step 1: Check direction to see if data deviate from H_0 in the direction specified by H_a

(If $\mu_1 < \mu_2$ then we expect t_s to be negative, if $\mu_1 > \mu_2$ then we expect t_s to be positive.)

a. If no, then p-value > 0.5

b. If yes, then proceed to step 2

■ Step 2: The p-value of the data is the one-tailed area beyond t_s

Slide 24 Stat 13, UCLA, Jon Dineen

One Tailed t Tests

Example: Cholesterol (cont')

RECALL: Group 1 = Medication, Group 2 = Placebo

Suppose it is reasonable to assume that $\mu_1 < \mu_2$, in other words the researcher is hoping to show that this new medication lowers cholesterol. The appropriate hypotheses would be

$$H_0: \mu_1 - \mu_2 = 0$$

$$H_a: \mu_1 - \mu_2 < 0$$

Calculate the p-value for this test

Step 1: t_s was calculated as -0.716. Check that the data deviate in the direction of H_a .


Slide 25

Stat 13, UCLA, Ivo Dinov


One Tailed t Tests

Step2: using 17 df, the p-value > 0.20

$p > 0.20$



What would the p-value be if the researchers had specified an upper tailed hypothesis?



Slide 26

Stat 13, UCLA, Ivo Dinov

One Tailed t Tests

Example: Soil pH (cont')

Suppose researchers had reason to believe that the soil pH for Location 1 was greater than Location 2.

Two-Sample T-Test and CI: Location 1, Location 2

Two-sample T for Location 1 vs Location 2

	N	Mean	StDev	SE Mean
Location 1	9	7.932	0.100	0.033
Location 2	9	7.460	0.222	0.074

$$\text{Difference} = \mu_1 - \mu_2$$

Estimate for difference: 0.472222

95% lower bound for difference: 0.326361

T-Test of difference = 0 (vs >): T-Value = 5.81

P-Value = 0.000 DF = 11

Slide 27

Stat 13, UCLA, Ivo Dinov

One Tailed t Tests

- P-values for a directional alternative are 1/2 of a non-directional

- assuming the direction matched H_a

- It is easier to reject H_0 with a one-tailed alternative

- However it is important that we decide on the direction of H_a before the data is collected

- If the data do not match the direction of H_a we conclude that the data do not indicate that the true means differ

- However t_s may be statistically significant in the other tail

- In this case we would want to look for methodological errors in the experiment

Slide 28

Stat 13, UCLA, Ivo Dinov

The Wilcoxon-Mann-Whitney

- Also known as the **rank sum test**

- http://www.socr.ucla.edu/htmls/SOCR_Analyses.html

- <http://socr.stat.ucla.edu/Applets.dir/WilcoxonRankSumTable.html>

- This hypothesis test is **also used to compare two independent samples**

- This procedure is different from the independent t test because it is **valid even if the population distributions are not normal**

- In other words, we can use this test as a fair substitute when we cannot meet the required normality assumption of the t test

- WMW is called **distribution-free** or non-parametric test

- This test doesn't focus on a parameter like the mean, instead it examines the distributions of the two groups

Slide 29

Stat 13, UCLA, Ivo Dinov

The Wilcoxon-Mann-Whitney

- Keep in mind that this is another hypothesis test, so there are still four major parts to consider

- #1 The hypotheses:

- H_0 : The population distributions of Y_1 and Y_2 are the same

- H_a : The population distributions of Y_1 and Y_2 are the different

- This could also be directional: distribution of Y_1 is less than Y_2 ; OR distribution of Y_1 is greater than Y_2

- #2 The test statistic:

- denoted by U_s

- measures the degree of separation between the two samples

- a large value of U_s indicates that the two samples are well separated with little overlap

- a small value of U_s indicates that the two samples are not well separated with much overlap

Slide 30

Stat 13, UCLA, Ivo Dinov

The Wilcoxon-Mann-Whitney

- #3 The p-value:
 - <http://socr.stat.ucla.edu/Applets.dir/WilcoxonRankSumTable.html>
 - http://www.socr.ucla.edu/htmls/SOCR_Analyses.html
 - Critical Values are also in table 6 on p.680
 - Method very similar to using the t table
 - find the appropriate row and then search for a number closest to the test statistic
 - don't need to worry about doubling the p-value for a two-tailed test (assuming we go to the right row header)
- #4 Conclusion:
 - Similar to the conclusion of an independent t test, but not linked to any parameter (for example the difference in means)

Slide 31 Stat 13, UCLA, Ivo Dinov

The Wilcoxon-Mann-Whitney


- The Method:
 - Step 1: Arrange the data in increasing order
 - Step 2: Determine K_1 and K_2
 - K_1 : for each observation in group 1, count the number of observations in the second group that are smaller. Use 1/2 for tied observations. K_1 is the sum of these ranks.
 - K_2 : for each observation in group 2, count the number of observations in the first group that are smaller. Use 1/2 for tied observations. K_2 is the sum of these ranks.
 - CHECK: if you have done the procedure correctly $K_1 + K_2 = n_1 n_2$
 - Step 3: If the test is non-directional then U_s is the larger of K_1 and K_2 . If the test is directional then U_s is the K that jives with the direction of H_a (if H_a is $Y_1 > Y_2$ then $U_s = K_1$, if H_a is $Y_1 < Y_2$ then $U_s = K_2$)
 - Step 4: Determine the critical value
 - $n =$ larger of n_1 and n_2
 - $n' =$ smaller of n_1 and n_2
 - Step 5: Bracket the p-value

Slide 32 Stat 13, UCLA, Ivo Dinov

The Wilcoxon-Mann-Whitney

Example: The urinary fluoride concentration (ppm) was measured both for a sample of livestock grazing in an area previously exposed to fluoride pollution and also for a similar sample of livestock grazing in an unpolluted area.

Polluted	Unpolluted
21.3	10.1
18.7	18.3
21.4	17.2
17.1	18.4
11.1	20.0
20.9	
19.7	




Does the data suggest that the fluoride concentration for livestock grazing in the polluted region is larger than that for the unpolluted region? Test using $\alpha = 0.01$.

Slide 33 Stat 13, UCLA, Ivo Dinov

The Wilcoxon-Mann-Whitney

Check Normality:



Slide 34 Stat 13, UCLA, Ivo Dinov

The Wilcoxon-Mann-Whitney

#1 The hypotheses:

H_0 : urinary fluoride values do not differ between the polluted and unpolluted regions.

H_a : the polluted region has a higher livestock urinary fluoride than the unpolluted region.

#2 The test statistic:

For this we need to deploy the WMW method shown a few slides earlier.

Slide 35 Stat 13, UCLA, Ivo Dinov

The Wilcoxon-Mann-Whitney

Let Polluted be group 1, and Unpolluted be group 2

Step 1: arrange the data in increasing order

# Unpolluted Below	Polluted $N_1=7$	Unpolluted $N_2=5$	# Polluted Below
1	11.1	10.1	0
1	17.1	17.2	2
4	18.7	18.3	2
4	19.7	18.4	2
5	20.9	20.0	4
5	21.3		
5	21.4		

$K_1 = 25$ $K_2 = 10$

Step 2: Determine K_1 and K_2

CHECK: $25 + 10 = 35 = (7)(5)$

Slide 36 Stat 13, UCLA, Ivo Dinov

The Wilcoxon-Mann-Whitney

Step 3: H_a : Polluted (Y_1) > Unpolluted (Y_2) so U_s is K_1

$$K_1 = 25 = U_s$$

$$K_2 = 10$$

Step 4:

$$n = 7$$

$$n' = 5$$

$$\alpha = 0.01$$

#3 The p-value:

$$0.2 > p > 0.1$$

Slide 37 Stat 13, UCLA, Ivo Dinno

The Wilcoxon-Mann-Whitney

#4 CONCLUSION: These data show that the [urinary fluoride concentration ppm for livestock grazing in polluted region is not greater than in the unpolluted region](#) ($P > 0.1$)

NOTE: No mention of the population means!

Corresponding Minitab output:

Mann-Whitney Test and CI: Polluted, Unpolluted

	N	Median
Polluted	7	19.700
Unpolluted	5	18.300

Test statistic is calculated using a different formula than our text, but W is the test statistic from the output

Point estimate for ETA1-ETA2 is 1.400
96.5 Percent CI for ETA1-ETA2 is (-2.897, 8.602)

W = 53.0
Test of ETA1 = ETA2 vs ETA1 > ETA2 is significant at 0.1278

The p-value is calculated using the computer, but is not labeled well.

Slide 38 Stat 13, UCLA, Ivo Dinno

The Wilcoxon-Mann-Whitney

Practice:

Say $n = 7$, $n' = 5$ and $U_s = 32$

Two-tailed p-value: $0.01 < p < 0.02$

One-tailed p-value: $0.005 < p < 0.01$

Say $n = 7$, $n' = 5$ and $U_s = 36$

Impossible, for these sample sizes U_s cannot be larger than 35!

Slide 39 Stat 13, UCLA, Ivo Dinno

The Wilcoxon-Mann-Whitney



- Why does this procedure make sense?

Suppose $n_1 = 3$ and $n_2 = 2$

$$K_1 + K_2 = (3)(2) = 6$$

we know that $K_1 + K_2$ should sum to 6

The relative magnitudes of K_1 and K_2 indicate the overlap in Y_1 and Y_2



Slide 40 Stat 13, UCLA, Ivo Dinno

The Wilcoxon-Mann-Whitney

- Conditions for the WMW:
 - Data are from random samples
 - Observations are independent
 - Samples are independent
- Remember: normality will not matter for this test

Slide 41 Stat 13, UCLA, Ivo Dinno

Wilcoxon-Mann-Whitney vs. Independent Test

- Both answer the same question, but treat data differently.
 - W-M-W uses rank ordering
 - Positive: doesn't depend on normality or population parameters
 - Negative: distribution free lacks power because it doesn't use all the info in the data
 - T-test uses actual Y values
 - Positive: Incorporates all of the data into calculations
 - Negative: Must meet normality assumption
 - neither is superior
 - If your data are normally distributed use the T-test
 - If your data are not normal use the WMW test

- http://www.socr.ucla.edu/htmls/SOCR_Analyses.html

Slide 42 Stat 13, UCLA, Ivo Dinno