

## Homework 1. Stat 202a. Due Tue, Oct 9, 10:30am.

You must work on the homework INDEPENDENTLY! Collaborating on this homework will be considered cheating. Submit your homework to me by email to [stat202a@stat.ucla.edu](mailto:stat202a@stat.ucla.edu). Your homework solution should be a single PDF document. The first pages should be your *output* from the problems above. This preferably means you typing out your answers and explaining them briefly in a clear, readable way, rather than simply cutting and pasting your R output. After that, on subsequent pages, include all your *code* for these problems. Your functions `pi2` and `pi3`, for example, should be included in the code section.

### 1. Approximating $\pi$ .

a) Write a function called `pi2(n)` that approximates  $\pi$  as a function of  $n$ , using the approximation  $\pi = \lim_{n \rightarrow \infty} \sqrt{6 \sum_{k=1}^n k^{-2}}$ . Evaluate `pi2(10j)` for  $j = 0, 1, 2, \dots, 6$ .

b) Write a function `pi3(n)` that approximates  $\pi$  as a function of  $n$ , by simulating random points in the square with vertices  $(-1, -1)$ ,  $(-1, 1)$ ,  $(1, 1)$ , and  $(1, -1)$ , seeing what fraction of them are in the unit circle [the circle with radius 1 centered at the origin], and then converting this fraction into an estimate of  $\pi$ . Evaluate `pi3(10j)` for  $j = 0, 1, 2, \dots, 6$ . For  $j=6$ , plot your simulated points, using different plotting symbols for simulated points inside and outside the unit circle. There is no need for you to plot the unit circle also.

### 2. Approximating $e$ .

a) Write a function `e2(n)` that approximates Euler's number,  $e \sim 2.71828$ , as a function of  $n$ . There are various ways to do this; choose one and explain which one you chose. Evaluate `e2(10j)` for  $j = 0, 1, 2, \dots, 6$ .

b) Plot `e2(n)` as a function of  $n$ , from  $n = 1$  to 100,000. (You do not need to plot every value of  $n$ . Just plot  $n=10^j$  for  $j = 0, 1, 2, \dots, 6$ .)

### 3. Assessing estimates of the 90th percentile of 100 iid uniform(0,1) random variables.

The R function `quantile()` implements a somewhat complex interpolation method in order to estimate a particular quantile, such as the 90th percentile. We will compare the estimate in `quantile()` with simpler estimates.

a) Write a function that takes as input a vector of length 100 and outputs the 90th of the 100 values sorted from smallest to largest. Note that the input vector might not be sorted.

b) Write a function to find the 91st of the sorted vector of 100 values.

c) Write a function that outputs the average of the 90th and 91st of the sorted vector of 100 values.

d) For each of your functions in parts a-c, as well as the function `quantile(x, 0.9)`, do the following:

(i) Generate 100 iid uniform (0,1) random variables, and calculate your estimate of the 90th percentile.

(ii) Repeat step (i) 100,000 times.

(iii) Plot the sample mean of the first  $m$  of your estimates, as a function of  $m$ . (continued on next page)

e) Report the ultimate sample mean of your 100,000 estimates, for each of the four estimates. Which of the 4 estimates appears to be the best?