

### Homework 3. Stat 202a. Due Thu, Nov 12, 11am.

You must work on the homework INDEPENDENTLY! Collaborating on this homework will be considered cheating. Submit your homework by email to [stat202a@stat.ucla.edu](mailto:stat202a@stat.ucla.edu). **Late homeworks will not be accepted!** Your homework solution should be a single PDF document. The first pages should be your *output* from the problems above. After that, on subsequent pages, include all your *code* for these problems.

#### 1. Approximation of an infinite series in C.

It is well known that  $1 - 1/2 + 1/3 - 1/4 + 1/5 - 1/6 + \dots = \ln(2)$ .

Write a C function called *alt2(n)* that computes the first  $n$  terms in this series, as a function of  $n$ . Call your C function from R to evaluate *alt2(n)* for various  $n$ . Using R, plot *alt2(n)* vs.  $n$ , for  $n$  ranging from some small number up to 1 million. You may set up your range of the y-axis in a way that you feel is appropriate. You do not need to show *alt2(n)* for all values of  $n$  and should not plot *alt2* for very small values of  $n$  if they are off the plot.

#### 2. Kernel density estimation in C and plotted in R.

Write a C function to compute a Gaussian kernel density estimate for univariate data. The inputs to the function should be two integers,  $m$  and  $n$ , a vector  $g$  of  $m$  gridpoints at which to calculate the estimates, a vector  $x$  consisting of the  $n$  observed data points, and a vector  $y$  of length  $m$  which will contain the resulting density estimates.

Gather data on all earthquakes of magnitude at least 3.0 in the longitude range -122.0 to -118.0 and latitude range 34.0 to 38.0, from Jan 1, 1960 to Jan 1 2018, from [http://service.scedc.caltech.edu/eq-catalogs/date\\_mag\\_loc.php](http://service.scedc.caltech.edu/eq-catalogs/date_mag_loc.php). Input the data into R. Use minimum magnitude = 3.0, maximum magnitude = 9.0, min depth = -5km, max depth = 100km, event type = earthquake, geographic type = local. There should be 2472 events. Take this vector of earthquake magnitudes, and use your C function to make a kernel density estimate of the earthquake magnitudes, using a Gaussian kernel with bandwidth selected using the rule of thumb suggested by Scott (1992). You may calculate this bandwidth in R. Let  $\{m_1, m_2, \dots, m_{100}\}$  = a vector of 100 equally spaced magnitudes spanning the observed range of magnitudes in your dataset, compute your kernel estimates on this grid using the C function, and plot your kernel density estimates  $\hat{f}(m_1)$ ,  $\hat{f}(m_2)$ , ...,  $\hat{f}(m_{100})$ .

**Output:** Your output for this assignment should be a pdf document containing the following, in this order.

Figure 1. A plot of *alt2(n)* versus  $n$ , for  $n$  ranging up to 1 million.

Figure 2. A plot of your kernel density estimates  $\hat{f}(m_1)$ ,  $\hat{f}(m_2)$ , ...,  $\hat{f}(m_{100})$  versus  $m$ .

After these 2 figures, include all of your C code, followed by all of your R code.