

University of California, Los Angeles
Department of Statistics

Statistics 100B

Instructor: Nicolas Christou

The F distribution

Definition:

Let $U \sim \chi_{n_1}^2$ and $V \sim \chi_{n_2}^2$. If U and V are independent the ratio

$\frac{\frac{U}{n_1}}{\frac{V}{n_2}}$ follows the F distribution with numerator d.f. n_1 and denominator d.f. n_2 .

We write $X \sim F_{n_1, n_2}$.

The probability density function of $X \sim F_{n_1, n_2}$ is:

$$f(x) = \frac{\Gamma(\frac{n_1+n_2}{2})}{\Gamma(\frac{n_1}{2})\Gamma(\frac{n_2}{2})} \left(\frac{n_1}{n_2}\right)^{\frac{n_1}{2}} x^{\frac{n_1}{2}-1} \left(1 + \frac{n_1}{n_2}x\right)^{-\frac{1}{2}(n_1+n_2)}, \quad 0 < x < \infty.$$

Proof:

Let $X \sim \frac{\frac{U}{n_1}}{\frac{V}{n_2}}$ and $W = V$. Solving for U we get $U = \frac{n_1}{n_2}XW$. Since U, V are independent the joint pdf of U and V is

$$f(u, v) = \frac{u^{\frac{n_1}{2}-1} e^{-\frac{u}{2}}}{\Gamma(\frac{n_1}{2}) 2^{\frac{n_1}{2}}} \times \frac{v^{\frac{n_2}{2}-1} e^{-\frac{v}{2}}}{\Gamma(\frac{n_2}{2}) 2^{\frac{n_2}{2}}}$$

The inverse of the Jacobian of the transformation is $\frac{n_1}{n_2}W$ (why?). Therefore the joint pdf of X and W is

$$f(x, w) = \frac{1}{\Gamma(\frac{n_1}{2})\Gamma(\frac{n_2}{2}) 2^{\frac{(n_1+n_2)}{2}}} \left(\frac{n_1 x w}{n_2}\right)^{\frac{n_1}{2}-1} w^{\frac{n_2}{2}-1} e^{-\frac{w}{2} \left(\frac{n_1 x}{n_2} + 1\right)} \frac{n_1 w}{n_2},$$

or

$$f(x, w) = \frac{\left(\frac{n_1}{n_2}\right)^{\frac{n_1}{2}} x^{\frac{n_1}{2}-1}}{\Gamma(\frac{n_1}{2})\Gamma(\frac{n_2}{2}) 2^{\frac{(n_1+n_2)}{2}}} w^{\frac{n_1+n_2}{2}-1} e^{-\frac{w}{2} \left(\frac{n_1 x}{n_2} + 1\right)}$$

Finally to find the marginal pdf of X we integrate the joint $f(x, w)$ w.r.t. w :

$$f(x) = \int_0^\infty \frac{\left(\frac{n_1}{n_2}\right)^{\frac{n_1}{2}} x^{\frac{n_1}{2}-1}}{\Gamma(\frac{n_1}{2})\Gamma(\frac{n_2}{2}) 2^{\frac{(n_1+n_2)}{2}}} w^{\frac{n_1+n_2}{2}-1} e^{-\frac{w}{2} \left(\frac{n_1 x}{n_2} + 1\right)} dw$$

Change the variable of integration by using the transformation, $y = \frac{w}{2} \left(\frac{n_1 x}{n_2} + 1\right)$. It follows that $dw = \frac{2}{\frac{n_1}{n_2}x+1} dy$ and $w = \frac{2y}{\frac{n_1}{n_2}x+1}$. Therefore,

$$f(x) = \int_0^\infty \frac{\left(\frac{n_1}{n_2}\right)^{\frac{n_1}{2}} x^{\frac{n_1}{2}-1}}{\Gamma(\frac{n_1}{2})\Gamma(\frac{n_2}{2}) 2^{\frac{(n_1+n_2)}{2}}} \left(\frac{2y}{\frac{n_1}{n_2}x+1}\right)^{\frac{n_1+n_2}{2}-1} e^{-y} \left(\frac{2}{\frac{n_1}{n_2}x+1}\right) dy$$

This is simplified to

$$f(x) = \frac{\Gamma(\frac{n_1+n_2}{2})}{\Gamma(\frac{n_1}{2})\Gamma(\frac{n_2}{2})} \left(\frac{n_1}{n_2}\right)^{\frac{n_1}{2}} x^{\frac{n_1}{2}-1} \left(1 + \frac{n_1}{n_2}x\right)^{-\frac{1}{2}(n_1+n_2)}, \quad \text{why?}$$