

100A

Ying Nian Wu

Discrete

Continuous

STATS 100A: RANDOM VARIABLES

Ying Nian Wu

Department of Statistics
University of California, Los Angeles

Some pictures are taken from the internet.
Credits belong to original authors.





Random variables

100A

Ying Nian Wu

Discrete

Continuous

Connection to events:

Randomly sample a person ω from a population Ω .



$X(\omega)$: gender of ω , $\Omega \rightarrow \{0, 1\}$.

$Y(\omega)$: height of ω , $\Omega \rightarrow \mathbb{R}^+$.

$A = \{\omega : X(\omega) = 1\}$. $P(A) = P(X = 1)$. Discrete.

$B = \{\omega : Y(\omega) > 6\}$. $P(B) = P(Y > 6)$. Continuous.

We shall study random variables more systematically.

$\omega \in \Omega$ equally likely, but $X(\omega)$ and $Y(\omega)$ are not necessarily equally likely.





Discrete random variable

100A

Ying Nian Wu

Discrete

Continuous

Randomly sample a person ω from a population Ω of N people.



$X(\omega)$: eye color of person ω , $\Omega \rightarrow \{1(\text{blue}), 2(\text{brown}), 3, 4\}$.

$N(x)$ = number of people with eye color x .

Probability mass function, probability distribution, law:

$$X \sim p(x) = P(X = x) = \frac{N(x)}{N}.$$

x	1	2	3	4
$p(x)$	$p(1)$	$p(2)$	$p(3)$	$p(4)$
number	$N(1)$	$N(2)$	$N(3)$	$N(4)$





Discrete random variable


100A

Ying Nian Wu

Discrete

Continuous

Randomly sample a person ω from a population Ω of N people.



$X(\omega)$: number of siblings of person ω , $\Omega \rightarrow \{0, 1, 2, \dots\}$.

$N(x)$ = number of people with x siblings.

Probability mass function, probability distribution, law:

$$X \sim p(x) = P(X = x) = \frac{N(x)}{N}.$$

x	0	1	2	3	...
$p(x)$	$p(0)$	$p(1)$	$p(2)$	$p(3)$...
number	$N(0)$	$N(1)$	$N(2)$	$N(3)$...






Population average

100A

Ying Nian Wu

Discrete

Continuous



$N(x)$ = number of people with x siblings.

$$X \sim p(x) = P(X = x) = \frac{N(x)}{N}.$$

x	0	1	2	3	...
$p(x)$	$p(0)$	$p(1)$	$p(2)$	$p(3)$...
number	$N(0)$	$N(1)$	$N(2)$	$N(3)$...

Population average:

$$\mathbb{E}(X) = \frac{1}{N} \sum_{\omega \in \Omega} X(\omega) = \frac{1}{N} \sum_x xN(x) = \sum_x x \frac{N(x)}{N} = \sum_x xp(x).$$





Long run average: N^n reasoning


100A

Ying Nian Wu

Discrete

Continuous

Randomly sample a person ω from a population Ω of N people.
Each person ω carries a number $X(\omega)$.



Population average:

$$\mu = \mathbb{E}(X) = \sum_x xp(x).$$

Repeat random sampling n times independently \rightarrow

N^n equally likely sequences: Ω_n .

\bar{X} (sequence) = average of the sequence.

$\bar{X} \rightarrow \mu$ in probability as $n \rightarrow \infty$.






Law of large number: N^n reasoning

100A

Ying Nian Wu

Discrete

Continuous



Repeat random sampling n times independently \rightarrow
 N^n equally likely sequences: Ω_n .

\bar{X} (sequence) = average of the sequence.

$A = \{\text{sequence} : |\bar{X}(\text{sequence}) - \mu| \leq .01\}$: representative sequences.

$P(A) = \frac{|A|}{N^n} \rightarrow 1$ as $n \rightarrow \infty$.






Die rolling

100A

Ying Nian Wu

Discrete

Continuous



Randomly throw a point into $[0, 1]$, which bin $(1, 2, \dots, 6)$ it falls into?

$\omega \in \Omega = [0, 1]$, population of points (tiny balls), equally likely.
 $X(\omega)$ is the bin that ω belongs to, not necessarily equally likely.
Probability mass function, probability distribution, law:

$$X \sim p(x) = P(X = x) = \text{length of bin } x.$$

Population average (each point or tiny ball ω carries a number $X(\omega)$).

$$\mathbb{E}(X) = \sum_x xp(x).$$





Long run average

100A

Ying Nian Wu

Discrete

Continuous

Independent repetitions:

$p(x)$: how often $X = x$ in the long run (e.g., throw 1 million points into $[0, 1]$).

x	1	2	3	4	5	6
$p(x)$	0.1	0.1	0.2	0.2	0.1	0.3

x	1	2	3	4	5	6
#	0.1m	0.1m	0.2m	0.2m	0.1m	0.3m
%	10%	10%	20%	20%	10%	30%

$$\text{average} = \frac{(1 \times 0.1m + 2 \times 0.1m + 3 \times 0.2m + 4 \times 0.2m + 5 \times 0.1m + 6 \times 0.3m)}{1m}$$

$$\mathbb{E}(X) = \sum_x xp(x).$$





Expectation of function

100A

Ying Nian Wu

Discrete

Continuous

Function of random variable

x	1	2	3	4	5	6
$p(x)$	0.1	0.1	0.2	0.2	0.1	0.3

x	1	2	3	4	5	6
#	0.1m	0.1m	0.2m	0.2m	0.1m	0.3m
%	10%	10%	20%	20%	10%	30%

$$\text{average} = \frac{(1 \times 0.1m + 2 \times 0.1m + 3 \times 0.2m + 4 \times 0.2m + 5 \times 0.1m + 6 \times 0.3m)}{1m}$$

x	1	2	3	4	5	6	x
payoff	-\$30	-\$20	\$0	\$20	\$30	\$100	$h(x)$
	$h(1)$	$h(2)$	$h(3)$	$h(4)$	$h(5)$	$h(6)$	

$$\text{longrun average} = (-\$30) \times 0.1 + (-\$20) \times 0.1 + (\$0) \times 0.2 + (\$20) \times 0.2 + (\$30) \times 0.1 + (\$100) \times 0.3$$

$$\mathbb{E}(h(X)) = \sum_x h(x)p(x).$$





Utility

100A

Ying Nian Wu

Discrete

Continuous

Utility, reward, value

$$\mathbb{E}(h(X)) = \sum_x h(x)p(x).$$

Offer 1	
x	\$100
p(x)	1

$$E(X) = (\$100) \times 1 = \$100$$

Offer 2		
x	\$0	\$200
p(x)	1/2	1/2

$$E(X) = (\$0) \times \frac{1}{2} + (\$200) \times \frac{1}{2} = \$100$$

x: face value	\$0	\$100	\$200
h(x): perceived value	\$0	\$100	\$150

Offer 1: $\mathbb{E}[h(X)] = \$100 \times 1 = \$100.$

Offer 2: $\mathbb{E}[h(X)] = \$0 \times \frac{1}{2} + \$150 \times \frac{1}{2} = \$75.$





Variance

100A

Ying Nian Wu

Discrete

Continuous

$$\mathbb{E}(X) = \sum_x xp(x) = \mu (= \$0 \times 1/2 + \$200 \times 1/2 = \$100)$$

$$\begin{aligned}\text{Var}(X) &= \mathbb{E}[(X - \mu)^2] = \sum_x (x - \mu)^2 p(x) = \sigma^2 \\ &= (\$0 - \$100)^2 \times 1/2 + (\$200 - \$100)^2 \times 1/2 \\ &= \$^2 10,000.\end{aligned}$$

Long run average of squared deviation from the mean.

$$SD(X) = \sqrt{\text{Var}(X)} = \sigma (= \$100).$$

Extent of variation from the mean.





Variance

100A

Ying Nian Wu

Discrete

Continuous

x	1	2	3	4	5	6	x
payoff	-\$30	-\$20	\$0	\$20	\$30	\$100	$h(x)$
	$h(1)$	$h(2)$	$h(3)$	$h(4)$	$h(5)$	$h(6)$	

$$\text{longrun average} = (-\$30) \times 0.1 + (-\$20) \times 0.1 + (\$0) \times 0.2 + (\$20) \times 0.2 + (\$30) \times 0.1 + (\$100) \times 0.3$$

$$\mathbb{E}(h(X)) = \sum_x h(x)p(x).$$

$$\text{Var}[h(X)] = \mathbb{E}[(h(X) - \mathbb{E}(h(X)))^2].$$





Data

100A

Ying Nian Wu

Discrete

Continuous

$$\mathbb{E}(X) = \sum_x xp(x) = \mu.$$

$$\text{Var}(X) = \mathbb{E}[(X - \mu)^2] = \sum_x (x - \mu)^2 p(x) = \sigma^2.$$

Long run average of squared deviation from the mean.

Sampling $p(x) \rightarrow x_1, \dots, x_i, \dots, x_n$

(e.g., rolling a die $\rightarrow 2, 1, 6, 5, 3, 2, 5, 4, 3, \dots$)

$$\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i \rightarrow \mathbb{E}(X) = \mu.$$

$$s^2 = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2 \rightarrow \text{Var}(X) = \sigma^2$$





Linear transformation


100A

Ying Nian Wu

Discrete

Continuous

$$\mathbb{E}(h(X)) = \sum_x h(x)p(x).$$
$$Y = aX + b.$$



$$\begin{aligned}\mathbb{E}(Y) &= \mathbb{E}(aX + b) = \sum_x (ax + b)p(x) \\ &= \sum_x axp(x) + \sum_x bp(x) \\ &= a \sum_x xp(x) + b \sum_x p(x) = a\mathbb{E}(X) + b.\end{aligned}$$





Data

100A

Ying Nian Wu

Discrete

Continuous

Sampling $p(x) \rightarrow x_1, \dots, x_i, \dots, x_n$
(e.g., rolling a die $\rightarrow 2, 1, 6, 5, 3, 2, 5, 4, 3, \dots$)

$$y_i = ax_i + b.$$

$$\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i \rightarrow \mathbb{E}(X) = \mu.$$

$$\bar{y} = \frac{1}{n} \sum_{i=1}^n y_i = \frac{1}{n} \sum_{i=1}^n (ax_i + b) = a \frac{1}{n} \sum_{i=1}^n x_i + b = a\bar{x} + b.$$





Linear transformation

100A

Ying Nian Wu

Discrete

Continuous

$$\text{Var}(h(X)) = \mathbb{E}[(h(X) - \mathbb{E}(h(X)))^2].$$

$$Y = aX + b.$$

$$\mathbb{E}(Y) = a\mathbb{E}(X) + b.$$

$$\text{Var}(Y) = \mathbb{E}[(Y - \mathbb{E}(Y))^2].$$

$$\begin{aligned}\text{Var}(aX + b) &= \mathbb{E}[((aX + b) - \mathbb{E}(aX + b))^2] \\ &= \mathbb{E}[(aX + b - (a\mathbb{E}(X) + b))^2] \\ &= \mathbb{E}[(a(X - \mathbb{E}(X)))^2] \\ &= a^2\mathbb{E}[(X - \mathbb{E}(X))^2] = a^2\text{Var}(X).\end{aligned}$$





Data

100A

Ying Nian Wu

Discrete

Continuous

Sampling $p(x) \rightarrow x_1, \dots, x_i, \dots, x_n$
(e.g., rolling a die $\rightarrow 2, 1, 6, 5, 3, 2, 5, 4, 3, \dots$)

$$\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i \rightarrow \mathbb{E}(X) = \mu.$$

$$y_i = ax_i + b.$$

$$\bar{y} = \frac{1}{n} \sum_{i=1}^n (ax_i + b) = a \frac{1}{n} \sum_{i=1}^n x_i + b = a\bar{x} + b.$$

$$\frac{1}{n} \sum_{i=1}^n (y_i - \bar{y})^2 = \frac{1}{n} \sum_{i=1}^n (ax_i + b - (a\bar{x} + b))^2 = \frac{1}{n} \sum_{i=1}^n a^2 (x_i - \bar{x})^2.$$





Short-cut for variance

100A

Ying Nian Wu

Discrete

Continuous

$$\mu = \mathbb{E}(X).$$

$$\begin{aligned}\text{Var}(X) &= \mathbb{E}[(X - \mu)^2] \\ &= \mathbb{E}[X^2 - 2\mu X + \mu^2] \\ &= \mathbb{E}(X^2) - 2\mu\mathbb{E}(X) + \mu^2 \\ &= \mathbb{E}(X^2) - \mu^2 = \mathbb{E}(X^2) - [\mathbb{E}(X)]^2.\end{aligned}$$

$$\begin{aligned}\mathbb{E}[h(X) + g(X)] &= \sum_x [h(x) + g(x)]p(x) \\ &= \sum_x h(x)p(x) + \sum_x g(x)p(x) \\ &= \mathbb{E}[h(X)] + \mathbb{E}[g(X)].\end{aligned}$$





Bernoulli

100A

Ying Nian Wu

Discrete

Continuous

Flip a coin (probability of head is p)

$Z \sim \text{Bernoulli}(p)$

$Z \in \{0, 1\}$, $P(Z = 1) = p$ and $P(Z = 0) = 1 - p$.

$$\mathbb{E}(Z) = 0 \times (1 - p) + 1 \times p = p.$$

$$\begin{aligned}\text{Var}(Z) &= (0 - p)^2 \times (1 - p) + (1 - p)^2 \times p \\ &= p(1 - p)[p + (1 - p)] = p(1 - p).\end{aligned}$$

$$\mathbb{E}(Z^2) = p.$$

$$\text{Var}(Z) = \mathbb{E}(Z^2) - \mathbb{E}(Z)^2 = p - p^2 = p(1 - p).$$





Binomial

100A

Ying Nian Wu

Discrete

Continuous

Flip a coin (probability of head is p) n times independently.

X = number of heads.

$X \sim \text{Binomial}(n, p)$

$$P(X = k) = \binom{n}{k} p^k (1 - p)^{n-k}.$$

$\binom{n}{k}$ is the number of sequences with exactly k heads.

$p^k (1 - p)^{n-k}$ is the probability of each sequence with k heads.

e.g., $n = 3$,

$P(X = 2) = P(HHT) + P(HTH) + P(THH) = 3p^2(1 - p).$

$p = 1/2$, we have $P(X = k) = \binom{n}{k} / 2^n.$





Recall independence

100A

Ying Nian Wu

Discrete

Continuous


Definition 1:

$$P(A|B) = P(A).$$

Definition 2:

$$P(A \cap B) = P(A)P(B).$$

	M	F
College degree	20	20
No college degree		
	50	50






Binomial formula

100A

Ying Nian Wu

Discrete

Continuous



$$(H + T)^n = \sum_{k=0}^n \binom{n}{k} H^k T^{n-k}.$$

$$n = 1, H + T.$$

$$n = 2, (H + T)(H + T) = HH + HT + TH + TT.$$

$$n = 3, \text{ above } \times (H + T) =$$

$$HHH + HHT + HTH + HTT + THH + THT + TTH + TTT.$$





Binomial and Bernoulli

100A

Ying Nian Wu

Discrete

Continuous

$$X = Z_1 + Z_2 + \dots + Z_n,$$

where $Z_i \sim \text{Bernoulli}(p)$ independently.

$$\mathbb{E}(X) = \sum_{i=1}^n \mathbb{E}(Z_i) = np.$$

Due to independence of Z_i , $i = 1, \dots, n$,

$$\text{Var}(X) = \sum_{i=1}^n \text{Var}(Z_i) = np(1 - p).$$





Frequency

100A

Ying Nian Wu

Discrete

Continuous

X/n is the frequency of heads.

$$\mathbb{E}(X/n) = \mathbb{E}(X)/n = p.$$

$$\text{Var}(X/n) = \text{Var}(X)/n^2 = p(1-p)/n.$$

$\text{Var}(X/n) \rightarrow 0$ as $n \rightarrow \infty$.

$X/n \rightarrow p$, in probability

Law of large number

Probability = long run frequency





Recall survey sampling


100A

Ying Nian Wu

Discrete

Continuous

Population of N people, M males.
Repeat random sampling n times independently



→ N^n equally likely sequences.

For a sequence ω , $X(\omega)$ = number of males in ω .

$A_m = \{\omega : X(\omega) = m\}$: sequences with m males.

$|A_m| = \binom{n}{m} M^m (N - M)^{n-m}$. n blanks. Choose m blanks for males, the rest $n - m$ blanks for females. Each male blank has M choices. Each female blank has $N - M$ choices.





Survey sampling

100A

Ying Nian Wu

Discrete

Continuous

Population of N people. M males.

Sample a person, $p = M/N = \text{Prob}(\text{male})$.

$$\begin{aligned}P(A_m) &= P(X = m) = \frac{|A_m|}{|\Omega_n|} \\ &= \frac{\binom{n}{m} M^m (N - M)^{n-m}}{N^n} \\ &= \binom{n}{m} p^m (1 - p)^{n-m}.\end{aligned}$$

Most sequences are representative, $X/n \approx M/N = p$.





Binomial distribution

100A

Ying Nian Wu


Discrete

Continuous

Flip a coin n times independently, $p =$ probability of head.

$$p(x) = P(X = x) = \binom{n}{x} p^x (1-p)^{n-x}$$
$$x = 0, 1, \dots, n.$$

$p(x)$: probability mass function, probability distribution.



Survey sampling, poll before election, $p = M/N$.
Monte Carlo, $p = \pi/4$.





Law of large number: N^n reasoning


100A

Ying Nian Wu

Discrete

Continuous

Survey sampling, poll before election, $p = M/N$.



Among all N^n sequences in the hyper-population of sequences Ω_n , let

$$A = \left\{ \omega : \left| \frac{X(\omega)}{n} - p \right| \leq .01 \right\}.$$

consist of representative sequences.

$$P(A) = \frac{|A|}{|\Omega_n|} = \sum_{x \in [n(p-.01), n(p+.01)]} p(x) \rightarrow 1,$$

$X/n \rightarrow p$ in probability.

$\mathbb{E}(X/n) = p$: average of $X(\omega)/n$ in Ω_n .

$\text{Var}(X/n) = p(1-p)/n \rightarrow 0$: variance of $X(\omega)/n$ in Ω_n .





Binomial expectation

100A

Ying Nian Wu

Discrete

Continuous

$$\begin{aligned}\mathbb{E}(X) &= \sum_{x=0}^n xp(x) = \sum_{k=0}^n kP(X = k) \\ &= \sum_{k=0}^n k \frac{n!}{k!(n-k)!} p^k (1-p)^{n-k} \\ &= \sum_{k=1}^n np \frac{(n-1)!}{(k-1)!(n-k)!} p^{k-1} (1-p)^{n-k} \\ &= \sum_{k'=0}^{n'} np \binom{n'}{k'} p^{k'} (1-p)^{n'-k'} = np.\end{aligned}$$

$$k' = k - 1; n' = n - 1.$$





Binomial variance

100A

Ying Nian Wu

Discrete

Continuous

$$\begin{aligned}\mathbb{E}(X(X-1)) &= \sum_{x=0}^n x(x-1)p(x) = \sum_{k=0}^n k(k-1)P(X=k) \\ &= \sum_{k=0}^n k(k-1) \frac{n!}{k!(n-k)!} p^k (1-p)^{n-k} \\ &= \sum_{k=2}^n n(n-1)p^2 \frac{(n-2)!}{(k-2)!(n-k)!} p^{k-2} (1-p)^{n-k} \\ &= \sum_{k'=0}^{n'} n(n-1)p^2 \binom{n'}{k'} p^{k'} (1-p)^{n'-k'} \\ &= n(n-1)p^2.\end{aligned}$$

$$k' = k - 2; \quad n' = n - 2.$$





Binomial variance

100A

Ying Nian Wu

Discrete

Continuous

$$\mathbb{E}(X) = np.$$

$$\mathbb{E}(X(X - 1)) = \mathbb{E}(X^2) - \mathbb{E}(X) = n(n - 1)p^2.$$

$$\begin{aligned}\text{Var}(X) &= \mathbb{E}(X^2) - \mathbb{E}(X)^2 \\ &= n(n - 1)p^2 + np - (np)^2 \\ &= np - np^2 = np(1 - p).\end{aligned}$$





Geometric

100A

Ying Nian Wu

Discrete

Continuous

$T \sim \text{Geometric}(p)$

T is the number of flips to get the first head, if we flip a coin independently and the probability of getting a head in each flip is p .

$$P(T = k) = (1 - p)^{k-1}p.$$

e.g., $T = 1, H$

$T = 2, TH.$

$T = 3, TTH.$

$T = 4, TTTH.$

Waiting time.





Geometric expectation

100A

Ying Nian Wu

Discrete

Continuous

$T \sim \text{Geometric}(p)$

$$\begin{aligned}\mathbb{E}(T) &= \sum_{k=1}^{\infty} kP(T = k) \\ &= \sum_{k=1}^{\infty} kq^{k-1}p = p \sum_{k=1}^{\infty} \frac{d}{dq} q^k \\ &= p \frac{d}{dq} \sum_{k=1}^{\infty} q^k = p \frac{d}{dq} \left(\frac{1}{1-q} - 1 \right) \\ &= p \frac{1}{(1-q)^2} = \frac{1}{p}.\end{aligned}$$





Geometric series

100A

Ying Nian Wu

Discrete

Continuous

$$\begin{aligned}(1 - a)(1 + a + \dots + a^m) &= 1 + a + \dots + a^m \\ &\quad - (a + a^2 + \dots + a^m + a^{m+1}) \\ &= 1 - a^{m+1}.\end{aligned}$$

$$1 + a + \dots + a^m = \frac{1 - a^{m+1}}{1 - a}.$$

If $|a| < 1$,

$a^{m+1} \rightarrow 0$, as $m \rightarrow \infty$.






Quantum bit

100A

Ying Nian Wu

Discrete

Continuous



state vector = $\alpha|0\rangle + \beta|1\rangle$.

state vector rotates over time.

squared length = $|\alpha|^2 + |\beta|^2 = 1$ under rotation.

observer: $p(0) = |\alpha|^2$, $p(1) = |\beta|^2$.

$$\frac{1}{\sqrt{2}}|\text{cat}\rangle + \frac{1}{\sqrt{2}}|\text{dog}\rangle$$

Schrodinger cat: $P(\text{alive}) = (1/\sqrt{2})^2 = 1/2$.





Continuous random variable: density

100A

Ying Nian Wu


Discrete

Continuous

Randomly sample a person ω from a population Ω of N people.

$X(\omega)$: height of person ω .

Density or distribution of the N points.



$N(x)$ = number of people in $(x, x + \Delta x)$ (6 ft, 6 ft 1 inch),
precision = 1 inch.

Probability density function, probability distribution, law:

$$X \sim f(x) = \frac{P(X \in (x, x + \Delta x))}{\Delta x} = \frac{N(x)/N}{\Delta x}.$$

Mathematical idealization: $N \approx \infty$.





Population scatterplot and histogram

100A

Ying Nian Wu

Discrete

Continuous



Discretize x -axis into equally spaced bins $(x, x + \Delta x)$, e.g., (6 ft, 6 ft 1 inch), precision = 1 inch.

$$P(X \in (x, x + \Delta x)) = \frac{N(x)}{N} = f(x)\Delta x.$$

$f(x)$: height of bin $(x, x + \Delta x)$. $f(x)\Delta x$: area.

$$\sum_x \frac{N(x)}{N} = \sum_x f(x)\Delta x \rightarrow \int f(x)dx = 1.$$






Region under curve

100A

Ying Nian Wu

Discrete

Continuous



Randomly throw a point ω into the region Ω below curve $f(x)$.
 Ω : population of points (tiny squares or balls).
Let $X = X(\omega)$ be the horizontal coordinate of point ω .

$$P(X \in (x, x + \Delta x)) = f(x)\Delta x.$$

$$P(X \in (a, b)) = \sum_{x \in (a, b)} f(x)\Delta x \rightarrow \int_a^b f(x)dx.$$






Independent repetitions, sample scatterplot and histogram

100A

Ying Nian Wu

Discrete

Continuous



Repeat n times, collapse to x -axis, histogram.

N^n reasoning:

sample scatterplot (random) \approx population scatterplot (fixed),

sample histogram (random) \approx population histogram (fixed).





Point cloud

100A

Ying Nian Wu

Discrete



Continuous

Electron orbits around nucleus: wrong conception

Electron cloud, probability density function, $f(x)$

Wave function $\psi(x)$, evolves over time.

Observer: $f(x) = |\psi(x)|^2$.





Population



100A

Ying Nian Wu

Discrete

Continuous

Electron cloud, heat map, prob density



Population of N equally likely possibilities.

Mathematical idealization: $N \approx \infty$.

Prob density = prob mass in the cell / volume of cell.

Observer: $f(x) = |\psi(x)|^2$.






Cumulative density function

100A

Ying Nian Wu

Discrete

Continuous



$$F(x) = P(X \leq x) = \int_{-\infty}^x f(x)dx.$$

SAT score $x \rightarrow$ percentile $F(x)$.

Percentage of people below x .






Area and slope

100A

Ying Nian Wu

Discrete

Continuous



Area:

$$F(x + \Delta x) - F(x) = f(x)\Delta x.$$

Slope:

$$F'(x) = \frac{F(x + \Delta x) - F(x)}{\Delta x} = f(x).$$

Notation:

$$F'(x) = \frac{dF(x)}{dx} = \frac{d}{dx}F(x) = f(x).$$

$$dF(x) = F'(x)dx = f(x)dx$$





Expectation

100A

Ying Nian Wu

Discrete

Continuous


Recall discrete

$$P(X = x) = p(x).$$

$$\mathbb{E}(X) = \sum xP(X = x) = \sum xp(x).$$

Continuous

$$P(X \in (x, x + \Delta x)) = f(x)\Delta x.$$



$$\mathbb{E}(X) = \sum xP(X \in (x, x + \Delta x)) = \sum xf(x)\Delta x \rightarrow \int xf(x)dx.$$

Population average, long run average, center.





Population average

100A


Ying Nian Wu

Discrete

Continuous

Population Ω of N people. $X(\omega)$.

$N(x)$: number of people in $(x, x + \Delta x)$.



$$\begin{aligned}\mathbb{E}(X) &= \frac{1}{N} \sum_{\omega} X(\omega) = \frac{1}{N} \sum_x x N(x) \\ &= \sum_x x \frac{N(x)}{N} = \sum_x x P(X \in (x, x + \Delta x)) \\ &= \sum_x x f(x) \Delta x \rightarrow \int x f(x) dx.\end{aligned}$$



Expectation of function

100A

Ying Nian Wu

Discrete

Continuous


Recall discrete

$$P(X = x) = p(x).$$

$$\mathbb{E}[h(X)] = \sum h(x)P(X = x) = \sum h(x)p(x).$$

Continuous

$$P(X \in (x, x + \Delta x)) = f(x)\Delta x.$$



$$\begin{aligned}\mathbb{E}[h(X)] &= \sum h(x)P(X \in (x, x + \Delta x)) \\ &= \sum h(x)f(x)\Delta x \rightarrow \int h(x)f(x)dx.\end{aligned}$$






Data, long run average

100A

Ying Nian Wu

Discrete

Continuous



$$f(x) \rightarrow x_1, \dots, x_i, \dots, x_n.$$

$n(x)$ = number of points in $(x, x + \Delta x)$.

$$\begin{aligned} \bar{x} &= \frac{1}{n} \sum_{i=1}^n x_i = \frac{1}{n} \sum_x x n(x) = \sum_x x \frac{n(x)}{n} \\ &\rightarrow \sum x P(X \in (x, x + \Delta x)) \\ &= \sum x f(x) \Delta x \rightarrow \int x f(x) dx = \mathbb{E}(X). \end{aligned}$$

Same logic for $\mathbb{E}(h(X))$.





Variance

100A


Ying Nian Wu

Discrete

Continuous

Continuous

$$P(X \in (x, x + \Delta x)) = f(x)\Delta x.$$



$$\mathbb{E}(X) = \int x f(x) dx = \mu.$$

$$\text{Var}(X) = \mathbb{E}[(X - \mu)^2] = \int (x - \mu)^2 f(x) dx.$$

$$\text{Var}[h(X)] = \mathbb{E}[(h(X) - \mathbb{E}(h(X)))^2].$$

Fluctuation, volatility, spread.





Uniform




100A

Ying Nian Wu

Discrete

Continuous

$U \sim \text{Uniform}[0, 1]$, i.e., the density of U is
 $f(u) = 1$ for $u \in [0, 1]$ (or $f(u) = 1/(b - a)$ if $u \in [a, b]$),
 $f(u) = 0$ otherwise.



$$P(U \in (u, u + \Delta u)) = f(u)\Delta u = \Delta u.$$

Imagine 1 million points distributed uniformly in $[0, 1]$.

Number of points in $(u, u + \Delta u)$ is Δu million.

e.g., Number of points in $(.3, .31)$ is .01 million.





Uniform

100A

Ying Nian Wu

Discrete

Continuous

$$F(u) = P(U \leq u) = \begin{cases} 0 & 0 < u \\ u & 0 \leq u \leq 1 \\ 1 & u > 1 \end{cases}$$

$F(u)$: proportion of points below u .

$$\mathbb{E}(U) = \int_0^1 u f(u) du = \frac{1}{2}.$$

$$\mathbb{E}(U^2) = \int_0^1 u^2 f(u) du = \frac{1}{3}.$$

$$\text{Var}(U) = \mathbb{E}(U^2) - (\mathbb{E}(U))^2 = \frac{1}{3} - \frac{1}{4} = \frac{1}{12}.$$





Pseudo-random number generator

100A

Ying Nian Wu

Discrete

Continuous

Linear congruential method

Start from an integer X_0 , and iterate

$$X_{t+1} = aX_t + b \bmod M.$$

Output $U_t = X_t/M$. e.g., $a = 7^5$, $b = 0$, and $M = 2^{31} - 1$.
mod: divide and take the remainder, e.g., $7 = 2 \bmod 5$.

e.g., $a = 7$, $b = 1$, $M = 5$, $X_0 = 1$, then

$$X_1 = 1 \times 7 + 1 \bmod 5 = 3.$$

$$X_2 = 3 \times 7 + 1 \bmod 5 = 2.$$






Exponential

100A

Ying Nian Wu

Discrete

Continuous



$$T \sim \text{Exponential}(\lambda),$$

$$f(t) = \lambda e^{-\lambda t} \text{ for } t \geq 0,$$

$$f(t) = 0 \text{ for } t < 0.$$

$$P(T \in (t, t + \Delta t)) = \lambda e^{-\lambda t} \Delta t.$$

Imagine 1 million particles, mark the times when they decay.
 1 million points on real line. Their distribution is exponential.
 Number of points in $(t, t + \Delta t)$ is $\lambda e^{-\lambda t} \Delta t$ million.






Exponential

100A

Ying Nian Wu

Discrete

Continuous



$$\begin{aligned}
 F(t) &= \int_0^t f(t)dt = \int_0^t \lambda e^{-\lambda t} dt \\
 &= -e^{-\lambda t} \Big|_0^t = 1 - e^{-\lambda t}.
 \end{aligned}$$

$F(t)$: proportion of points below t

Half-life: $F(t_{\text{half}}) = P(T \leq t_{\text{half}}) = 1/2$.

1 million particles, by half life, half million will have decayed.





Exponential expectation

100A

Ying Nian Wu

Discrete

Continuous

$$\begin{aligned}\mathbb{E}(T) &= \int_0^{\infty} t\lambda e^{-\lambda t} dt \\ &= - \int_0^{\infty} t de^{-\lambda t} \\ &= -(te^{-\lambda t}|_0^{\infty} - \int_0^{\infty} e^{-\lambda t} dt) \\ &= -(0 - 0 + \frac{1}{\lambda} e^{-\lambda t}|_0^{\infty}) = \frac{1}{\lambda}.\end{aligned}$$






Integral by parts

100A

Ying Nian Wu

Discrete

Continuous



$$\frac{d}{dx}u(x)v(x) = u'(x)v(x) + u(x)v'(x).$$

$$d(uv) = u dv + v du.$$

$$\int [u'(x)v(x) + u(x)v'(x)] dx = u(x)v(x).$$

$$\int u(x)v'(x) dx = u(x)v(x) - \int v(x)u'(x) dx.$$

$$\int u dv = uv - \int v du.$$






Integral by parts

100A

Ying Nian Wu

Discrete

Continuous



$$\int u dv = uv - \int v du.$$

$$\int u(x)v'(x)dx = u(x)v(x) - \int v(x)u'(x)dx.$$

$$\frac{du(x)}{dx} = \frac{d}{dx}u(x) = u'(x); \quad du(x) = u'(x)dx.$$





Normal or Gaussian

100A




Ying Nian Wu

Discrete

Continuous

Let $Z \sim N(0, 1)$, i.e., the density of Z is

$$f(z) = \frac{1}{\sqrt{2\pi}} e^{-z^2/2}.$$



$$\int_{-2}^2 f(z) dz = 95\%.$$





Normal expectation

100A

Ying Nian Wu

Discrete

Continuous

Let $Z \sim N(0, 1)$, i.e., the density of Z is

$$f(z) = \frac{1}{\sqrt{2\pi}} e^{-z^2/2}.$$

$$\begin{aligned} \mathbb{E}(Z) &= \int_{-\infty}^{\infty} z \frac{1}{\sqrt{2\pi}} e^{-\frac{z^2}{2}} dz \\ &= -\frac{1}{\sqrt{2\pi}} e^{-\frac{z^2}{2}} \Big|_{-\infty}^{\infty} \\ &= 0. \end{aligned}$$

The density is symmetric around 0.





Normal variance

100A

Ying Nian Wu

Discrete

Continuous

Let $Z \sim N(0, 1)$, i.e., the density of Z is

$$f(z) = \frac{1}{\sqrt{2\pi}} e^{-z^2/2}.$$

$$\begin{aligned}\mathbb{E}(Z^2) &= \int_{-\infty}^{\infty} z^2 \frac{1}{\sqrt{2\pi}} e^{-\frac{z^2}{2}} dz \\ &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} (-z) de^{-\frac{z^2}{2}} \\ &= \frac{1}{\sqrt{2\pi}} \left(-ze^{-\frac{z^2}{2}} \Big|_{-\infty}^{\infty} - \int_{-\infty}^{\infty} e^{-\frac{z^2}{2}} d(-z) \right) \\ &= \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} e^{-\frac{z^2}{2}} dz = 1.\end{aligned}$$

$$\text{Var}(Z) = \mathbb{E}(Z^2) - (\mathbb{E}(Z))^2 = 1.$$





Variance

100A

Ying Nian Wu

Discrete

Continuous

For $X \sim f(x)$, let $\mu = \mathbb{E}(X)$.

$$\begin{aligned}\text{Var}(X) &= \mathbb{E}[(X - \mu)^2] \\ &= \mathbb{E}[X^2 - 2\mu X + \mu^2] \\ &= \mathbb{E}(X^2) - 2\mu\mathbb{E}(X) + \mu^2 \\ &= \mathbb{E}(X^2) - (\mathbb{E}(X))^2.\end{aligned}$$

$$\begin{aligned}\mathbb{E}[r(X) + s(X)] &= \int [r(x) + s(x)]f(x)dx \\ &= \int r(x)f(x)dx + \int s(x)f(x)dx \\ &= \mathbb{E}[r(X)] + \mathbb{E}[s(X)].\end{aligned}$$





Linear transformation

100A

Ying Nian Wu

Discrete

Continuous

For $X \sim f(x)$. Let $Y = aX + b$.

$$\begin{aligned}\mathbb{E}(Y) = \mathbb{E}(aX + b) &= \int (ax + b)f(x)dx \\ &= a \int xf(x)dx + b \int f(x)dx \\ &= a\mathbb{E}(X) + b.\end{aligned}$$

$$\begin{aligned}\text{Var}(Y) = \text{Var}(aX + b) &= \mathbb{E}[((aX + b) - \mathbb{E}(aX + b))^2] \\ &= \mathbb{E}[(aX + b - (a\mathbb{E}(X) + b))^2] \\ &= \mathbb{E}[a^2(X - \mathbb{E}(X))^2] \\ &= a^2\mathbb{E}[(X - \mathbb{E}(X))^2] = a^2\text{Var}(X).\end{aligned}$$







Data

100A

Ying Nian Wu

Discrete

Continuous



Sampling $f(x) \rightarrow x_1, \dots, x_i, \dots, x_n$
(e.g., random number generator $\rightarrow .22, .31, .92, .45, \dots$)

$$\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i \rightarrow \mathbb{E}(X) = \mu.$$

$$s^2 = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2 \rightarrow \text{Var}(X) = \sigma^2$$



Data

100A

Ying Nian Wu

Discrete

Continuous

Sampling $f(x) \rightarrow x_1, \dots, x_i, \dots, x_n$
(e.g., random number generator $\rightarrow .22, .31, .92, .45, \dots$)

$$y_i = ax_i + b.$$

$$\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i \rightarrow \mathbb{E}(X) = \mu.$$

$$\bar{y} = \frac{1}{n} \sum_{i=1}^n y_i = \frac{1}{n} \sum_{i=1}^n (ax_i + b) = a \frac{1}{n} \sum_{i=1}^n x_i + b = a\bar{x} + b.$$

$$\frac{1}{n} \sum_{i=1}^n (y_i - \bar{y})^2 = \frac{1}{n} \sum_{i=1}^n (ax_i + b - (a\bar{x} + b))^2 = \frac{1}{n} \sum_{i=1}^n a^2 (x_i - \bar{x})^2.$$





Change of density under linear transformation

100A


Ying Nian Wu

Discrete

Continuous

Change of variable

$X \sim f(x)$, $Y = aX + b$ ($a > 0$). $Y \sim g(y)$.



$$y = ax + b, \quad x = (y - b)/a.$$

$$P(X \in (x, x + \Delta x)) = P(Y \in (y, y + \Delta y)).$$

$$f(x)\Delta x = g(y)\Delta y.$$

$$g(y) = f(x) \frac{\Delta x}{\Delta y} = f((y - b)/a) / a.$$

Space warping, stretching or squeezing.





Normal or Gaussian

100A

Ying Nian Wu

Discrete

Continuous

Let $Z \sim N(0, 1)$, i.e., the density of Z is

$$f(z) = \frac{1}{\sqrt{2\pi}} e^{-z^2/2}.$$

Let $X = \mu + \sigma Z$. $Z = (X - \mu)/\sigma$. Then

$$\mathbb{E}(X) = \mathbb{E}(\mu + \sigma Z) = \mu + \sigma \mathbb{E}(Z) = \mu.$$

$$\text{Var}(X) = \text{Var}(\mu + \sigma Z) = \sigma^2 \text{Var}(Z) = \sigma^2.$$

$$f(z)\Delta z = g(x)\Delta x.$$

$$\begin{aligned} g(x) &= f(z) \frac{\Delta z}{\Delta x} \\ &= f((x - \mu)/\sigma) / \sigma \\ &= \frac{1}{\sqrt{2\pi}\sigma^2} \exp\left[-\frac{(x - \mu)^2}{2\sigma^2}\right]. \end{aligned}$$





Normal or Gaussian

100A

Ying Nian Wu


Discrete

Continuous

Let $Z \sim N(0, 1)$. Let $X = \mu + \sigma Z$. $Z = (X - \mu)/\sigma$.
 $X \sim N(\mu, \sigma^2)$,

$$f(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp \left[-\frac{(x - \mu)^2}{2\sigma^2} \right].$$

(we now use $f(x)$ to denote the density of X .)



$$P(\mu - 2\sigma \leq X \leq \mu + 2\sigma) = P(-2 \leq Z \leq 2) = 95\%.$$